

Dear Parents/Guardians:

Missoula County Public Schools policy requires your consent in order to administer the over-the-counter medications (OTC) described below. All <u>other medications & treatments require an additional form with the signature of your child's</u> <u>health care provider</u>. (This includes all prescription, over the counter and complementary and alternative medicine (CAM)). Forms available at school and on the district website under the Health Services tab.

I give permission for the school nurse and/or other designee to administer the medications below to

tudents Name	Date of Birth	Grade
My child is allergic to		
My child has previously taken Tylenol (acetami	inophen) 📃 Yes 🗋	No
My child has previously taken Ibuprofen (Motr	in or Advil) 🛛 🗌 Yes 🗌	No
My child has a sensitivity to dyes in medication Advil). Yes No (If yes, you may be a	. ,	etaminophen (Tylenol) or ibuprofen (Motrin or

I understand that I need to supply the school with liquid or chewable medications or medications for field trips if needed. I understand school procedure is that I am to give the medication to the school office in its original container. School personnel will discard the medications supplied at the end of the school year if a parent does not pick them up prior to this.

Parent/ Guardian Signature

Date

## STANDING ORDERS FOR STUDENTS

	Acetaminophen (Tylenol) Dose	Ibuprofen (Advil/Motrin) Dose
Kindergarten	1 ½ teaspoon= 7.5 ml=240mg of liquid	1 ½ teaspoon= 7.5 ml= 150mg of liquid
	acetaminophen 160mg (5 ml) concentration	ibuprofen 100mg (5 ml) concentration
Grades 1-4	One 325 mg tablet OR one 500mg tablet (up	One 200 mg tablet or 2 teaspoons (10 ml)
	to 3 teaspoons (15 ml))	
Grades 5-8	Two 325 mg tablets OR ONE 500mg tablet	Two 200mg tablets or 4 teaspoons (20 ml)
	(3-4 teaspoons (15-20 ml))	
Frequency/ max	Up to every 4 hours, no more than 1300 mg	Once every 8 hours.
dose	in any 8 hour period	

- <u>Benadryl (diphenhydramine) 25mg</u> for minor allergic reaction to include swelling at site of sting, and/or hives or itching at area of contact of allergen. The school will notify the parent and school nurse of allergic reaction prior to medication administration when possible.
- For students age 12 and older only <u>Tums (calcium carbonate)</u> 1-2 tablets chewed, no more than twice per day for minor stomach distress.